The case for specifying the "ideal" target trial

Margarita Moreno-Betancur^{1,2,*}, Rushani Wijesuriya^{1,2}, John B. Carlin^{1,2}

¹ Clinical Epidemiology and Biostatistics Unit, Department of Paediatrics, University of Melbourne, Melbourne, Australia

² Clinical Epidemiology and Biostatistics Unit, Murdoch Children's Research Institute, Melbourne, Australia

*Corresponding author: margarita.moreno@unimelb.edu.au

21 May 2024

Abstract

The target trial is an increasingly popular conceptual device for guiding the design and analysis of

observational studies that seek to perform causal inference. As tends to occur with concepts like this,

there is variability in how certain aspects of the approach are understood, which may lead to

potentially consequential differences in how the approach is taught, implemented, and interpreted in

practice. In this paper, we consider two of these aspects: how the target trial should be specified, and

relatedly, how the target trial fits within a formal causal inference framework. We first describe two

challenges with what we call the "aligned" approach to target trial specification, which is common in

evaluations of medical interventions using healthcare databases and seeks to specify a target trial that

is closely aligned to the observational data so that all protocol components apart from randomization

can be closely emulated. We then argue how these challenges can be circumvented by an approach

that focusses on specifying the "ideal" target trial: a trial with certain idealized aspects that ensure it is

relevant to answer the research question. In essence, this approach is applicable in a broader range of

settings and enables systematic assessment of all potential sources of causal bias. Importantly, this

view provides clarification for how the target trial approach fits within a formal causal inference

framework.

Keywords: Target trial, causal inference, bias

1

Introduction

The target trial^{1–4} is an increasingly popular⁵ conceptual device for guiding the design and analysis of observational studies that seek to perform causal inference, which we understand as the task of addressing "what if" questions about the impact of interventions.⁶ The general idea is to specify the hypothetical randomized controlled trial that one would have liked to conduct to examine the causal effect of interest – the target trial – and then to perform the analysis of the observational data in a way that emulates the target trial as closely as possible.

As tends to occur once concepts like this are released into the wild, there is variability in how the approach is understood (see e.g. ⁷⁻¹²). Unfortunately, different perspectives lead to potentially consequential differences in how the approach is taught, implemented, and interpreted in practice. Key differences concern how the target trial should be specified, and relatedly, how the target trial fits within a formal causal inference framework ^{6,13-16}. It is widely understood that the specified target trial needs to be pragmatic ³ (e.g. not placebo-controlled or blinded) so that it is feasible to emulate it with observational data, but beyond this there is little guidance regarding the details of target trial specification.

In many applications (e.g. ^{17–22}), the approach has been to specify an "aligned" target trial: a trial that is closely aligned to the observational data in the sense that it ensures that all protocol components apart from randomization can be closely emulated. For example, in evaluations of pharmacological interventions using healthcare databases, the eligibility criteria may include being enrolled in the health care system to which the database corresponds, and the outcome measure in the target trial may be specified to be the same as that captured in the database. When it is feasible to specify an aligned target trial that is relevant to the research question, then by construction it can be closely emulated except for randomization, for which we must rely on the unverifiable assumption of no residual confounding given the selected measured confounders. In this context, it is said that a correct target trial emulation eliminates common sources of bias except possibly for confounding⁴.

The are two challenges with this "aligned" approach to target trial specification. Firstly, in some contexts it may be difficult or impossible to specify an aligned target trial that is relevant to the research question. In many epidemiologic studies of complex social, behavioral, and environmental exposures, there are necessary distinctions between a relevant target trial and its emulation beyond the lack of randomization, even if the data are drawn from high-quality longitudinal cohort studies or other studies with primary data collection. These distinctions and the potential biases emerging from them need to be explicitly acknowledged. Secondly, even when it is feasible to specify an aligned target trial and sufficient adjustment for confounding appears possible in the observational study, both the aligned target trial and its emulation remain subject to the same sources of bias that actual trials are subject to, a fact that is not made explicit with this approach.

In this paper, we illustrate the challenges with the aligned approach by way of an example in respiratory epidemiology. We then propose that researchers should instead (or in addition, where specifying the aligned trial is feasible) aim to specify the "ideal" target trial: a trial with certain idealized aspects that ensure that it is relevant to answer the research question. This approach is applicable to the broader range of contexts in which the target trial framework is already used (e.g. ^{23–25}) and enables systematic assessment of all potential sources of causal bias. Further, this view provides clarification for how the target trial framework fits within a formal causal inference framework.

Challenges with specifying an aligned target trial

Our illustrative example considers an investigation of the effect of breastfeeding on risk of asthma at age 6 years based on data from HealthNuts, a prospective longitudinal cohort study.²⁶ The original manuscript did not use the target trial framework, but in Table 1 we have proposed possible ideal and (somewhat) aligned target trials for the emulation implicit in (a simplified version of) that analysis.

The first challenge with the aligned approach to specifying the target trial is that close alignment (apart from randomization) may be infeasible. In the asthma example, in attempting to specify an aligned trial, we observe necessary misalignment across three protocol components (columns 3 and 4

of Table 1). In *Eligibility criteria*, the aligned trial would necessarily follow infants from birth, but in the cohort study infants are only followed up from 11-15 months of age. Furthermore, to be in the emulation study sample, families would have had to consent to participate in HealthNuts (a multifaceted long-term cohort study), whereas no such condition would be needed in the aligned trial. Finally, in the emulation, the analytic sample needs to be determined with consideration for missing data in variables such as the treatment (or adherence to it), confounders and eligibility criteria, but there would be no such missing data in the aligned trial. In *Treatment strategies*, the aligned trial would record the age at initiating formula or other foods in quasi-real time, whereas the emulation must rely on parent report at recruitment to the cohort study (when infant aged 11-15 months). In *Outcome*, the aligned trial would ascertain the asthma outcome through clinical diagnosis whereas the emulation must rely on parent-report and medication use data.

The described misalignments are potentially consequential in that they represent sources of causal bias beyond confounding that could arise in emulating it. For example, if there are unmeasured common causes of reaching age 11-15 months and consenting to participate in the cohort at this post-treatment time-point, the emulation could be subject to selection bias. Multivariable missing data are also an important source of bias depending on the causal structure^{27–30}.

The second challenge is that, even if we had a study where alignment were achievable (say, if the cohort study had followed children from birth, clinically diagnosed asthma, etc.), there would still be three potential sources of bias relative to the ultimate causal estimand of interest, defined by the ideal target trial (see below), but these remain tacit when only the aligned target trial and emulation are specified. These biases arise in the first two protocol components. In *Eligibility criteria*, selection bias may arise if the observational study sample is not representative of the true target population – so called type II selection or generalizability bias³¹. In the example, this could arise if the effect of breastfeeding on asthma differs by whether the parent can read and understand English, which is possible if there is effect modification by ethnicity, for which English fluency is a strong proxy.

Further selection bias may arise due to missing data in the outcome, or in predictors of either missingness in outcome, selection into the study sample or adherence to treatment. For instance, bias

can arise if there are unmeasured common causes of outcome and outcome missingness such as parental healthcare-seeking behavior. In *Treatment strategies*, confounding bias could arise in estimation of the per-protocol effect if there are unmeasured predictors of non-adherence and outcome.

The ideal target trial

An approach that focusses on specifying the ideal target trial avoids the two challenges above. Firstly, this approach would be automatically applicable to scenarios where a relevant target trial must meet idealized conditions that differ from the observational study. For instance, it would be natural to specify a gold-standard measure for the outcome in the ideal target trial, such as clinical diagnosis of asthma in the example. Secondly, unlike the aligned target trial, the ideal target trial captures what is ultimately the target causal effect of interest. For instance, in the example, findings of the study aim to inform public health guidelines for all neonates in Victoria who can be breastfed, not just those whose parents speak English (Table 1, column 2). As such, the ideal target trial is the referent that is needed to systematically assess all sources of causal bias. Table 2 summarizes the statistical issues that can arise in the ideal and aligned target trials and the emulation, and the full set of assumptions required for emulating the ideal target trial.

More broadly, the concept of the ideal target trial clarifies the relationship between the target trial framework and a formal causal inference framework 6,13–16. The Figure shows what can be conceived as the three essential steps in the formal approach to causal inference: 32 (1) (non-parametric) estimand definition, (2) (non-parametric) identification and (3) estimation. Asymptotically (i.e. if we imagine an infinite sample size), components A to E of the ideal target trial can be seen to fully operationalize what is arguably the causal estimand of ultimate interest⁶: in the example, it is the average causal effect of (fully adherent) exclusive breastfeeding in the first 6 months of life on asthma risk in the target population. In the ideal (but not the aligned) target trial, the intention-to-treat and per-protocol effects (component F), are identical to each other and are equal to this effect (see eAppendix). The definition of the target estimand entails important value judgements about the question that we seek to address, which can be referred to as estimand assumptions.³²

The process of designing the emulation of components A to E of the ideal target trial is the practical operationalization of the identification step, amounting to expressing the estimand as a function of the observable data. Each component is successfully emulated only if unverifiable causal identifiability assumptions underpinning each emulation decision hold, for example no error in selected measures or no unmeasured common causes of missingness and outcome given selected predictors of missingness (see Table 2 and eAppendix). If these assumptions are violated, causal biases (i.e. discrepancies relative to the target causal estimand due to study design aspects) would arise. While the *Statistical analysis plan* for the ideal target trial is trivial, that for the emulation must delineate the analytic approaches that will be used in light of those assumptions, for example to adjust for the selected confounders and predictors of missingness. Analytic approaches often rely on parametric modelling assumptions. The execution of this plan when analyzing the data corresponds to the estimation step and it is at this stage that one must reckon with finite sample size issues and potential non-causal biases (e.g. due to misspecification of parametric models).

Conclusion

There are important nuances in the implementation and interpretation of the target trial approach that arise as it gets used across different substantive areas. Given the accelerating uptake of the approach over recent years, examining these nuances is important for strengthening its use in practice. In this paper, we considered the question of how the target trial should be specified, and the related matter of how the approach fits within a formal causal inference question. We highlighted two challenges with an approach to target trial specification that aims for close alignment with the observational study apart from randomization of the intervention: first, in many settings specifying an aligned target trial is not feasible, and, second, even when feasible, the aligned trial may not fully operationalize the target estimand of ultimate interest. We have argued that aiming to specify the ideal target trial instead (or in addition, where the aligned version is feasible) broadens the applicability of the framework, provides a basis for systematic identification of all causal biases in a study and clarifies the connection with a formal causal inference framework. We believe our proposal is in line with

recommendations in a foundational target trial paper¹, which stated that one must provide an explanation of the way in which the [aligned] target trial is distinct from the ideal trial.

The appeal of the aligned approach to specifying the target trial is that it distinguishes issues specific to observational studies (confounding) from issues arising in both observational and randomized studies (non-generalizability, non-adherence, outcome missingness). This may help to improve understanding of observational study evidence among consumers of trial evidence. However, this agenda may conflict with that of strengthening causal inference more broadly. Failing to specify the ideal target trial, which defines the ultimate estimand, means losing sight of necessary assumptions and thus limitations and potential biases, which may result in overconfidence in evidence arising from target trial emulations that mirrors overconfidence in evidence arising from randomized trials³³.

In fact, specifying the ideal target trial can be argued to be important for actual randomized trials too, emphasizing the power of the target trial framework as a unifying concept for the analysis of both observational studies and trials⁶. Specifying the ideal trial is also fundamental in comparisons of multiple studies: multiple trials, multiple observational studies³⁴ or combinations of both, such as when benchmarking observational studies against an actual trial^{35,36}. Indeed, comparing one study with the other is the usual tendency whereas each study should be appraised against a common reference, defined in terms of the ultimate target estimand, as has been argued elsewhere.³⁴

To conclude, specification of the ideal target trial ensures a clear definition of the estimand, which is the critical first step in a logically coherent formal approach to causal inference. Taking this approach, the target trial framework operationalizes and clearly distinguishes the key concepts of estimand definition, identification and estimation,³² which we believe is essential for clear causal thinking in practice.

TABLES

Table 1. Ideal and aligned target trials, and emulation, for the illustrative example

Protocol	Ideal target trial	Aligned ^a target trial	Emulation
A. Eligibility criteria ^b	Study sample Representative sample of neonates born in Melbourne in 2007-2011 who can be breastfed	Study sample Representative sample of neonates born in Melbourne in 2007-2011 who can be breastfed with a parent/guardian who can read and understand English	Study sample Sample of infants aged 11-15 months in 2007-2011 participating in 170 council-run immunization sessions across Melbourne who can be breastfed with a parent/guardian who can read and understand English whose families consent to participate in a cohort study
	Analytic sample • All participants in study sample	Analytic sample • All participants in study sample regardless of missing data in outcome Approach to handling missing data and other potential sources of selection bias	Analytic sample • All participants in study sample regardless of missing data in exposure, outcome and confounders. Approach to handling missing data and other potential sources of selection bias
		requires identification of predictors of missingness & other selection processes	requires identification of predictors of missingness & other selection processes
B. Treatment strategies	 Exclusive breast-feeding during the first 6 months Non-exclusive or no breast-feeding during the first 6 months 	 Exclusive breast-feeding during the first 6 months Non-exclusive or no breast-feeding during the first 6 months 	 Exclusive breast-feeding during the first 6 months Non-exclusive or no breast-feeding during the first 6 months
		Measure of adherence: Quasi real-time measure of age at initiating formula or other foods	Measure of adherence: Based on 12-month parent report of age at initiating formula or other foods (see "Assignment procedures")
		Approach to non-adherence adjustment requires selection of predictors of adherence at baseline and post-baseline	Approach to non-adherence adjustment and to handle exposure measurement error requires selection of predictors of adherence at baseline and post-baseline, and predictor of misclassified exposure
C. Assignment procedures	Randomization at recruitment without blind assignment	Randomization at recruitment without blind assignment	No randomization, no blinding. Individuals assigned at birth to strategy consistent with the 12-month parent report of age at initiating formula or other foods. For individuals who cannot be unambiguously assigned to these strategies at time zero (i.e. those initiating exclusive breast-feeding at birth), the method of random allocation or cloning is used (Hernan, Robins et al. 2016) Grace periods could be used if relevant data were available
			Approach to baseline confounding adjustment requires selection of confounders at baseline
D. Follow-up period	Follow-upStarts: BirthEnds: Child aged 6 years	Follow-up • Starts: Birth • Ends: Child aged 6 years	Follow-up • Starts: Child aged 11-15 months • Ends: Child aged 6 years
E. Outcome	Outcome measure • Clinically diagnosed asthma at age 6, with systematic and blind ascertainment	Outcome measure • Clinically diagnosed asthma at age 6, with systematic and blind ascertainment	Outcome measure • Parental report of doctor-diagnosed asthma or use of common asthma medication in the previous 12 months
			Approach to handle outcome misclassification requires selection of predictors of misclassified outcome
F. Causal contrasts ^c	ITT _{ideal} PPE _{ideal}	ITT _{align} PPE _{align}	ITT _{obs} PPE _{obs}
G. Statistical analysis plan	N.B. ITT _{ideal} =PPE _{ideal} ITT _{ideal} , PPE _{ideal} : Compare proportions with asthma between treatment arms	ITT _{align} : Compare proportions with asthma between treatment arms, using an approach such as multiple imputation (MI) or inverse probability weighting (IPW) to handle missing outcome data	ITT _{obs} : Cannot be emulated in this case given treatment not unambiguously assigned at birth (Hernan, Sauer et al. 2016), but if this were not the case, we would compare proportions with asthma between treatment arms, using an approach like g-computation or IPW to adjust for baseline confounders, and MI or IPW to handle missing outcome data
		PPE _{align} : Compare proportions with asthma between treatments arms, using MI or IPW to handle missing outcome data and IPW to adjust for non-adherence (individuals who deviate from assigned strategy are censored)	PPE _{obs} : Compare proportions with asthma between treatment arms, using g-computation or IPW to adjust for baseline confounders, MI or IPW to handle multivariable missing data and IPW to adjust for non-adherence (individuals who deviate from assigned strategy are censored)
		N.B. No attempt to correct for a non- representative sample – stronger assumptions needed	N.B. No attempt to correct for a non- representative sample, or misclassified exposure and outcome – stronger assumptions needed

^a Aligned to the extent possible in this example, though still presents important and potentially consequential misalignments (see text).

^b Following Lu et al. 2022,³¹ the target population is the population about which inference is sought, the study sample is the population enrolled into the study (trial or cohort), and the analytic sample is the portion of the study sample used in the analysis.

 $^{^{}c} \ See \ eAppendix \ for \ formal \ definitions \ of \ ITT_{ideal} \ \ PPE_{ideal}, \ ITT_{align}, \ \ PPE_{align}, \ ITT_{obs} \ and \ \ PPE_{obs} \ in \ this \ example.$

Table 2. Statistical issues that can arise in ideal and aligned target trials and emulations, and full set of assumptions required for emulating the ideal target trial

Protocol component	Issue	Ideal target trial	Aligned target trial	Emulation	Assumption for emulating the ideal target trial ^a
A. Eligibility criteria	Non-representative study sample	No	Yes	Yes	No unmeasured predictors of selection that could induce bias
	Missing data	No	Yes; missing data in outcome, and predictors of adherence, missingness and selection	Yes; missing data in outcome, treatment, confounders, eligibility criteria and in predictors of adherence, missingness, selection and misclassification	No unmeasured predictors of missingness that could induce bias
B. Treatment strategies	Non-adherence	No	Yes	Yes	No unmeasured predictors of non-adherence that could induce bias
	Exposure measurement error	No	No	Yes	No unmeasured predictors of exposure measurement error that could induce bias
C. Assignment procedures	Confounding	No	No	Yes	No unmeasured confounders at baseline that could induce bias
D. Follow-up period	Misalignment between time zero, treatment assignment and eligibility	No	No	Yes	[This can lead to a non-representative study sample and/or measurement error in exposure – both issues covered above]
E. Outcome measure	Outcome measurement error	No	No	Yes	No unmeasured predictors of outcome measurement error that could induce bias

^a Whether unmeasured predictors can cause bias in each case depends on the causal diagram/structure, e.g. see Zhang et al. 2024²⁸ for the case of multivariable missingness.

FIGURE




Figure. Diagram depicting how the target trial framework fits within the three essential steps in the formal causal inference framework.

REFERENCES

- 1. Hernán MA, Robins JM. Using Big Data to Emulate a Target Trial When a Randomized Trial Is Not Available. *Am J Epidemiol*. 2016;183(8):758-764.
- Hernán MA, Sauer BC, Hernández-Díaz S, Platt R, Shrier I. Specifying a target trial prevents immortal time bias and other self-inflicted injuries in observational analyses. *J Clin Epidemiol*. 2016;79:70-75. doi:10.1016/j.jclinepi.2016.04.014
- 3. Hernán MA. Methods of Public Health Research Strengthening Causal Inference from Observational Data. *N Engl J Med.* 2021;385(15):1345-1348. doi:10.1056/nejmp2113319
- 4. Hernán MA, Wang W, Leaf DE. Target Trial Emulation: A Framework for Causal Inference From Observational Data. *JAMA*. 2022;328(24):2446-2447. doi:10.1001/JAMA.2022.21383
- Hansford HJ, Cashin AG, Jones MD, et al. Reporting of Observational Studies Explicitly
 Aiming to Emulate Randomized Trials: A Systematic Review. *JAMA Netw Open*.

 2023;6(9):E2336023. doi:10.1001/jamanetworkopen.2023.36023
- 6. Hernan MA, Robins J. Causal Inference: What If. Boca Raton: Chapman & Hall/CRC; 2020.
- Pearce N, Vandenbroucke JP. Are Target Trial Emulations the Gold Standard for Observational Studies? *Epidemiology*. 2023;34(5):614-618.
 doi:10.1097/EDE.000000000001636
- 8. Dang LE, Balzer LB. Start with the Target Trial Protocol, Then Follow the Roadmap for Causal Inference. *Epidemiology*. 2023;34(5):619-623. doi:10.1097/EDE.000000000001637
- Swanson SA. The Causal Effects of Causal Inference Pedagogy. *Epidemiology*.
 2023;34(5):611-613. doi:10.1097/EDE.000000000001639
- 10. Didelez V, Haug U, Garcia-Albeniz X. Re: Are Target Trial Emulations the Gold Standard for Observational Studies? *Epidemiology*. 2024;35(1):E3. doi:10.1097/EDE.0000000000001667
- 11. Pearce N, Vandenbroucke J. Are Target Trial Emulations the Gold Standard for Observational Studies? The authors respond. *Epidemiology*. 2024;35(2):E5-E6. doi:10.1097/EDE.000000000001668
- 12. De Stavola BL, Gomes M, Katsoulis M. Transparency and Rigor: Target Trial Emulation Aims to Achieve Both. *Epidemiology*. 2023;34(5):624-626.

- doi:10.1097/EDE.0000000000001638
- 13. Robins J. A new approach to causal inference in mortality studies with a sustained exposure period-application to control of the healthy worker survivor effect. *Math Model*. 1986;7(9-12):1393-1512. doi:10.1016/0270-0255(86)90088-6
- 14. Rubin D. B. Estimating causal effects of treatment in randomized and nonrandomized studies. *J Educ Psychol.* 1974;66(5):688-701.
- 15. Pearl J. Causality. New York: Cambridge University Press; 2000.
- 16. van der Laan MJ, Rose S. *Targeted Learning. Causal Inference for Observational and Experimental Data*. New York: Springer; 2011.
- 17. Hernández-Díaz S, Huybrechts KF, Chiu YH, Yland JJ, Bateman BT, Hernán MA. Emulating a Target Trial of Interventions Initiated During Pregnancy with Healthcare Databases: The Example of COVID-19 Vaccination. *Epidemiology*. 2023;34(2):238-246. doi:10.1097/EDE.0000000000001562
- Dickerman BA, García-Albéniz X, Logan RW, Denaxas S, Hernán MA. Avoidable flaws in observational analyses: an application to statins and cancer. *Nat Med*. 2019;25(10):1601-1606. doi:10.1038/s41591-019-0597-x
- García-Albéniz X, Hsu J, Hernán MA. The value of explicitly emulating a target trial when using real world evidence: an application to colorectal cancer screening. *Eur J Epidemiol*. 2017;32(6):495-500. doi:10.1007/s10654-017-0287-2
- Szmulewicz AG, Martínez-Alés G, Logan R, et al. Antipsychotic drugs in first-episode psychosis: A target trial emulation in the FEP-CAUSAL Collaboration. *Am J Epidemiol*. April 2024. doi:10.1093/AJE/KWAE029
- 21. Rein SM, Lodi S, Logan RW, et al. Integrase strand-transfer inhibitor use and cardiovascular events in adults with HIV: an emulation of target trials in the HIV-CAUSAL Collaboration and the Antiretroviral Therapy Cohort Collaboration. *Lancet HIV*. 2023;10(11):e723-e732. doi:10.1016/S2352-3018(23)00233-3
- Dickerman BA, García-Albéniz X, Logan RW, Denaxas S, Hernán MA. Evaluating Metformin
 Strategies for Cancer Prevention: A Target Trial Emulation Using Electronic Health Records.

- Epidemiology. 2023;34(5):690-699. doi:10.1097/EDE.000000000001626
- 23. Moreno-Betancur M, Lynch JW, Pilkington RM, et al. Emulating a target trial of intensive nurse home visiting in the policy-relevant population using linked administrative data. *Int J Epidemiol.* 2023;52(1):119-131. doi:10.1093/IJE/DYAC092
- Rojas-Saunero LP, Labrecque JA, Swanson SA. Invited Commentary: Conducting and Emulating Trials to Study Effects of Social Interventions. *Am J Epidemiol*. 2022;191(8):1453-1456. doi:10.1093/aje/kwac066
- Goldfeld S, Downes M, Gray S, et al. Household income supplements in early childhood to reduce inequities in children's development. *Soc Sci Med*. 2024;340:116430. doi:10.1016/J.SOCSCIMED.2023.116430
- 26. Peters RL, Kay T, McWilliam VL, et al. The Interplay Between Eczema and Breastfeeding Practices May Hide Breastfeeding's Protective Effect on Childhood Asthma. *J Allergy Clin Immunol Pract*. 2021;9(2):862-871.e5. doi:10.1016/j.jaip.2020.09.006
- Moreno-Betancur M, Lee KJ, Leacy FP, White IR, Simpson JA, Carlin JB. Canonical Causal Diagrams to Guide the Treatment of Missing Data in Epidemiologic Studies. *Am J Epidemiol*. 2018;187(12):2705-2715.
- Zhang J, Dashti SG, Carlin JB, Lee KJ, Moreno-Betancur M. Recoverability and estimation of causal effects under typical multivariable missingness mechanisms. *Biometrical J*. 2023;66(2200326).
- 29. Lee KJ, Carlin JB, Simpson JA, Moreno-Betancur M. Assumptions and analysis planning in studies with missing data in multiple variables: moving beyond the MCAR/MAR/MNAR classification. *Int J Epidemiol*. 2023:1268-1275. doi:10.1093/ije/dyad008
- 30. Tompsett D, Zylbersztejn A, Hardelid P, De Stavola B. Target Trial Emulation and Bias Through Missing Eligibility Data: An Application to a Study of Palivizumab for the Prevention of Hospitalization Due to Infant Respiratory Illness. Am J Epidemiol. 2023;192(4):600-611. doi:10.1093/aje/kwac202
- 31. Lu H, Cole SR, Howe CJ, Westreich D. Toward a Clearer Definition of Selection Bias When Estimating Causal Effects. *Epidemiology*. 2022;33(5):699-706.

- doi:10.1097/EDE.0000000000001516
- 32. Moreno-Betancur M. The target trial: A powerful device beyond well-defined interventions. *Epidemiology*. 2020;32(1):291-294.
- 33. Deaton A, Cartwright N. Understanding and misunderstanding randomized controlled trials. Soc Sci Med. 2018;210:2-21. doi:10.1016/j.socscimed.2017.12.005
- 34. Downes M, O'Connor M, Olsson CA, et al. Causal inference in multi-cohort studies using the target trial framework to identify and minimize sources of bias (pre-print).
 2024;arXiv:2206.11117.
- 35. Lodi S, Phillips A, Lundgren J, et al. Practice of Epidemiology Effect Estimates in Randomized Trials and Observational Studies: Comparing Apples With Apples. Am J Epidemiol. 2019;188(8):1569-1577. doi:10.1093/aje/kwz100
- 36. Dahabreh IJ, Robins JM, Hernán MA. Benchmarking Observational Methods by Comparing Randomized Trials and Their Emulations. *Epidemiology*. 2020;31(5):614-619. doi:10.1097/EDE.000000000001231

FUNDING: This work was supported by an Investigator Grant fellowship to MMB from the National Health and Medical Research Council [grant ID: 2009572]. The Murdoch Children's Research Institute is supported by the Victorian Government's Operational Infrastructure Support Program.

CONFLICTS OF INTEREST: None declared.

ACKNOWLEDGEMENTS: The authors would like to thank members of our local causal inference team for ongoing discussions surrounding these concepts. We would also like to thank Rachel Peters for providing details about the analysis of the cited breastfeeding and asthma publication that we used as the basis for the example.

eAPPENDIX

Formal definitions of causal contrasts for the example (cf. Table 1 of main text)

Let X denote a binary 0/1 indicator of treatment arm assignment and \overline{A} a vector of binary 1/0 indicators of received treatment each day over the first 6 months of the child's life (for each day, 1 indicates exclusive breastfeeding, 0 otherwise). Let Y denote the outcome, M_Y a binary 0/1 outcome missingness indicator, and we use superscripts to denote potential outcomes, e.g. $Y^{X=x}$ is the potential outcome when X is set to x. Let E_W denote an expectation over the covariate distribution W and denote by W_1 , W_2 and W_3 the covariate distributions in the study samples for the ideal target trial, aligned target trial and emulation, respectively.

In the ideal target trial, the causal contrasts in the mean difference scale are:

$$\mathsf{ITT}_{\mathsf{ideal}} \coloneqq E_{W_1}(Y^{X=1,M_Y=0}) - E_{W_1}(Y^{X=0,M_Y=0}) = E_{W_1}(Y^{X=1}) - E_{W_1}(Y^{X=0})$$

and

$$PPE_{ideal} := E_{W_1} (Y^{\bar{A} = \bar{1}, M_Y = 0}) - E_{W_1} (Y^{\bar{A} = \bar{0}, M_Y = 0}) = E_{W_1} (Y^{\bar{A} = \bar{1}}) - E_{W_1} (Y^{\bar{A} = \bar{0}}),$$

where ITT and PPE stand for intention-to-treat and per-protocol effect, respectively and the second equalities in each definition arise because there is no outcome missingness. Further, because there is no non-adherence, PPE_{ideal} = ITT_{ideal} and this is the ultimate causal estimand of interest: the average causal effect of (fully adherent) exclusive breastfeeding in the first 6 months of life on asthma risk in the target population. Further, the statistical analysis plan of the ideal target trial (Table 1 in main text) targets a function of the observable data that identifies (i.e. is equal to) this estimand. Indeed, $E_{W_1}(Y^{X=x}) = E_{W_1}(Y^{X=x}|X=x) = E_{W_1}(Y|X=x) \text{ in the ideal target trial.}$

In the aligned target trial, the causal contrasts are:

$$ITT_{align} := E_{W_2}(Y^{X=1,M_Y=0}) - E_{W_2}(Y^{X=0,M_Y=0})$$

and

$$PPE_{align} := E_{W_2} (Y^{\bar{A} = \bar{1}, M_Y = 0}) - E_{W_2} (Y^{\bar{A} = \bar{0}, M_Y = 0}).$$

Here, non-adherence means that $PPE_{align} \neq ITT_{align}$ in general, and whether the statistical analysis plan targets functions of the observable data that identify ITT_{align} and PPE_{align} respectively depends on unverifiable assumptions about the adherence and missingness mechanisms. In a typical case, the analyst might claim identifiability of ITT_{align} by assuming Y independent of M_Y given predictors of missingness used in an approach to handle missing data such as multiple imputation or inverse probability weighting. Given non-adherence, ITT_{align} is in general different from the ultimate causal estimand of interest, PPE_{ideal} (= ITT_{ideal}) and, given a discrepancy between W_1 and W_2 , so is PPE_{align} .

In the emulation, the observational analog of the intention-to-treat effect can be defined based on initiating exclusive breastfeeding in, say, the first day, week or month. Whatever the choice, denote this by $A_0 = 1$, and $A_0 = 0$ otherwise. Thus, in the emulation, the causal contrasts are:

$$ITT_{obs} := E_{W_3}(Y^{A_0 = 1, M_Y = 0}) - E_{W_3}(Y^{A_0 = 0, M_Y = 0})$$

and

$$PPE_{obs} := E_{W_3} (Y^{\bar{A} = \overline{1}, M_Y = 0}) - E_{W_3} (Y^{\bar{A} = \overline{0}, M_Y = 0}).$$

Whether the statistical analysis plan of the emulation targets functions of the observable data that identify ITT_{obs} and PPE_{obs} depends on an expanded set of unverifiable assumptions, about adherence, multivariable missingness, treatment receipt and measurement mechanisms. Given non-adherence, ITT_{obs} is in general different from the ultimate causal estimand of interest, PPE_{ideal} (= ITT_{ideal}) and, given a discrepancy between W_1 and W_3 , so is PPE_{obs} .